

# **Statistical Rethinking**

## **Winter 2019**

Lecture 09 / Week 5

### **Conditional Manatees**





# Stop testing, start thinking

- Off-the-shelf tools
  - $p$ -values
  - information criteria
  - linear models
- Good decisions benefit from **bespoke** tools
  - bespoke risk analysis
  - bespoke models

**bespoke** | bə'spōk |

adjective *chiefly British*

made for a particular customer or user: *a bespoke suit | bespoke kitchens | bespoke software systems | group tours and bespoke itineraries.*

- making or selling bespoke goods, especially clothing: *bespoke tailors.*

*Bespoke*



*Broke*



# *Leaders in New York and New Jersey Defend Shutdown for a Blizzard That Wasn't*

By MATT FLEGENHEIMER JAN. 27, 2015

Email

Share

Tweet

Save

More

It was an unprecedented step for what became, in New York City, a common storm: For the first time in its 110-year history, the subway system was shut down because of snow.

Transit workers, caught off guard by the shutdown that Gov. [Andrew M. Cuomo](#) announced on Monday, scrambled to grind the network to a halt within hours.


Residents moved quickly to find places



Mayor Bill de Blasio of New York City, with Sanitation Department workers in Manhattan on Monday, when he issued dire warnings about the storm. Yana Paskova for The New York Times



## World leader in global medium-range numerical weather prediction



# Blizzard calibration

- Was it bad to predict NYC blizzard from ECMWF?
- Other models were more accurate
- But welfare enhanced by being prepared => use extreme forecasts
- **Accuracy always matters, but it isn't all that matters**







# Fly



## *Armstrong Whitworth Whitley Mk V*

British Medium Bomber



Plastic model kit  
Plastik-Modellbausatz  
Plastikový model

### 1/72





AFRICA



CALISTA



DONNA



FLASH



FLOYD








Figure 8.1



rotor/wing

keel/engine

survival

Observe only:  
undamaged  
rotor/wing damaged

Figure 8.1



# Manatees and bombers


- *Conditioning*: Dependence on state
- Everything is conditional
  - On data
  - On model
  - On information state
- Influence of variable conditional on other variable(s)

# Interaction effects

- *Interactions*: Influence of predictor conditional on other predictor(s)
  - Influence of *sugar* in *coffee* depends on *stirring*
  - Influence of *gene* on *phenotype* depends on *environment*
  - Influence of *skin color* on *cancer* depends on *latitude*
- Generalized linear models (GLMs): All predictors interact to some degree
- Multilevel models: Massive interaction engines

# Interaction effects in DAGs

- In DAG, interaction doesn't look special:



- This just means:


$$coffee\ sweet = f(sugar, stirred)$$

- We have to figure out the function  $f$ .




*sugar*  $\longrightarrow$  *coffee sweet*  $\longleftarrow$  *stirred*

not an interaction



interaction





```
library(rethinking)  
data(rugged)  
d <- rugged
```





Figure 8.2



# The sermon on priors







Figure 8.3

# The value of being rugged

- Splitting the data is a bad idea:
  - No inference for how you split the data
  - Does not pool information
- How about adding a categorical variable for Africa?



# Category doesn't work

- Index variable for continent:

$$\mu_i = \alpha_{\text{CID}[i]} + \beta(r_i - \bar{r})$$

```
m8.4 <- quap(  
  alist(  
    log_gdp_std ~ dnorm( mu , sigma ) ,  
    mu <- a[cid] + b*( rugged_std - 0.215 ) ,  
    a[cid] ~ dnorm( 1 , 0.1 ) ,  
    b ~ dnorm( 0 , 0.3 ) ,  
    sigma ~ dexp( 1 )  
  ) ,  
  data=dd )
```




Figure 8.4



# Interaction

- Need to allow effect of ruggedness to depend upon continent

$$\mu_i = \alpha_{\text{CID}[i]} + \beta_{\text{CID}[i]}(r_i - \bar{r})$$

# Interaction

$$\mu_i = \alpha_{\text{CID}[i]} + \beta_{\text{CID}[i]}(r_i - \bar{r})$$

R code  
8.13

```
m8.5 <- quap(  
  alist(  
    log_gdp_std ~ dnorm( mu , sigma ) ,  
    mu <- a[cid] + b[cid]*( rugged_std - 0.215 ) ,  
    a[cid] ~ dnorm( 1 , 0.1 ) ,  
    b[cid] ~ dnorm( 0 , 0.3 ) ,  
    sigma ~ dexp( 1 )  
  ) ,  
  data=dd )
```

R code  
8.13

```
m8.5 <- quap(  
  alist(  
    log_gdp_std ~ dnorm( mu , sigma ) ,  
    mu <- a[cid] + b[cid]*( rugged_std - 0.215 ) ,  
    a[cid] ~ dnorm( 1 , 0.1 ) ,  
    b[cid] ~ dnorm( 0 , 0.3 ) ,  
    sigma ~ dexp( 1 )  
  ) ,  
  data=dd )
```

R code  
8.14

```
precis( m8.5 , depth=2 )
```

	mean	sd	5.5%	94.5%
a[1]	0.89	0.02	0.86	0.91
a[2]	1.05	0.01	1.03	1.07
b[1]	0.13	0.07	0.01	0.25
b[2]	-0.14	0.05	-0.23	-0.06
sigma	0.11	0.01	0.10	0.12







Figure 8.5

# Interpreting interactions

- Is hard
  - Add interaction => other parameters change meaning
  - Influence of predictor depends upon multiple parameters and their covariation

R code  
8.14

```
precis( m8.5 , depth=2 )
```

	mean	sd	5.5%	94.5%
a[1]	0.89	0.02	0.86	0.91
a[2]	1.05	0.01	1.03	1.07
b[1]	0.13	0.07	0.01	0.25
b[2]	-0.14	0.05	-0.23	-0.06
sigma	0.11	0.01	0.10	0.12

# Interactions are symmetric

- Effect of ruggedness depends upon continent:

$$\mu_i = \alpha_{\text{CID}[i]} + \beta_{\text{CID}[i]}(r_i - \bar{r})$$

- Effect of continent depends upon ruggedness:

$$\mu_i = \underbrace{(2 - \text{CID}_i)(\alpha_1 + \beta_1(r_i - \bar{r}))}_{\text{CID}[i]=1} + \underbrace{(\text{CID}_i - 1)(\alpha_2 + \beta_2(r_i - \bar{r}))}_{\text{CID}[i]=2}$$

# Interactions are symmetric





Figure 8.6

# Continuous interactions

- `data(tulips)`: 27 replicate blooms across three levels of both water and shade





# Tulip blooms

No interaction:

*water and shade have independent effects*

$$b_i \sim \text{Normal}(\mu_i, \sigma)$$

$$\mu_i = \alpha + \beta_w(w_i - \bar{w}) + \beta_s(s_i - \bar{s})$$

Interaction:

*water and shade have interdependent effects*

$$b_i \sim \text{Normal}(\mu_i, \sigma)$$

$$\mu_i = \alpha + \beta_w(w_i - \bar{w}) + \beta_s(s_i - \bar{s}) + \beta_{ws}(w_i - \bar{w})(s_i - \bar{s})$$



# How is interaction formed?

$$b_i \sim \text{Normal}(\mu_i, \sigma)$$

$$\mu_i = \alpha + \beta_w(w_i - \bar{w}) + \beta_s(s_i - \bar{s}) + \beta_{ws}(w_i - \bar{w})(s_i - \bar{s})$$

# How is interaction formed?

$$b_i \sim \text{Normal}(\mu_i, \sigma)$$

$$\mu_i = \alpha + \beta_w(w_i - \bar{w}) + \beta_s(s_i - \bar{s}) + \beta_{ws}(w_i - \bar{w})(s_i - \bar{s})$$

$$\mu_i = \alpha + \gamma_{w,i}W_i + \beta_sS_i$$

$$\gamma_{w,i} = \beta_w + \beta_{ws}S_i$$

# How is interaction formed?

$$b_i \sim \text{Normal}(\mu_i, \sigma)$$

$$\mu_i = \alpha + \beta_w(w_i - \bar{w}) + \beta_s(s_i - \bar{s}) + \beta_{ws}(w_i - \bar{w})(s_i - \bar{s})$$

$$\mu_i = \alpha + \gamma_{w,i}W_i + \beta_sS_i$$

$$\gamma_{w,i} = \beta_w + \beta_{ws}S_i$$

$$\mu_i = \alpha + \underbrace{(\beta_w + \beta_{ws}S_i)}_{\gamma_{w,i}} W_i + \beta_s S_i = \alpha + \beta_w W_i + \beta_s S_i + \beta_{ws} S_i W_i$$

# Tulip model – no interaction

```
m8.6 <- quap(  
  alist(  
    blooms_std ~ dnorm( mu , sigma ) ,  
    mu <- a + bw*water_cent + bs*shade_cent ,  
    a ~ dnorm( 0.5 , 0.25 ) ,  
    bw ~ dnorm( 0 , 0.25 ) ,  
    bs ~ dnorm( 0 , 0.25 ) ,  
    sigma ~ dexp( 1 )  
  ) ,  
  data=d )
```

R code  
8.23



# Plotting interaction

- Slope changes with values of other predictor, so use more than one plot
- Here, need three plots, *trptych*

trip•tych | 'triptik |

noun

a picture or relief carving on three panels, typically hinged together side by side and used as an altarpiece.  
• a set of three associated artistic, literary, or musical works intended to be appreciated together.




ORIGIN mid 18th cent. (denoting a set of three writing tablets hinged or tied together): from TRI-*'three,'* on the pattern of *diptych*.



Lewis Powell (1844–1865), before his hanging for conspiracy to assassinate Abraham Lincoln.

# Prior predictions

No interaction



# Posterior predictions

No interaction




Figure 8.7

# Tulip model – interaction


R code  
8.24

```
m8.7 <- quap(  
  alist(  
    blooms_std ~ dnorm( mu , sigma ) ,  
    mu <- a + bw*water_cent + bs*shade_cent + bws*water_cent*shade_cent ,  
    a ~ dnorm( 0.5 , 0.25 ) ,  
    bw ~ dnorm( 0 , 0.25 ) ,  
    bs ~ dnorm( 0 , 0.25 ) ,  
    bws ~ dnorm( 0 , 0.25 ) ,  
    sigma ~ dexp( 1 )  
  ) ,  
  data=d )
```

Interpreting parameters very hard! Plot.

# Prior predictions

No interaction



Interaction





Figure 8.8



# Posterior predictions

No interaction



Interaction







Figure 8.7

# Causal thinking

- Tulip experiment:



- Tulip reality:



# Interactions not always linear

- Suppose all tulip data collected under “cool” temperatures
- Under “hot” temperature, tulips do not bloom
- Interaction, but not a linear one
  - blooms goes to zero at threshold




# Higher order interactions

- Just keep multiplying:

$$\begin{aligned} y_i &\sim \text{Normal}(\mu_i, \sigma), \\ \mu_i &= \alpha + \boxed{\beta_1 x_{1i} + \beta_2 x_{2i} + \beta_3 x_{3i}} \quad \text{main effects} \\ &\quad + \boxed{\beta_{12} x_{1i} x_{2i} + \beta_{13} x_{1i} x_{3i} + \beta_{23} x_{2i} x_{3i}} \quad \text{2-way interactions} \\ &\quad + \boxed{\beta_{123} x_{1i} x_{2i} x_{3i}} \quad \text{3-way interaction} \end{aligned}$$

# Higher order interactions

- Dangers of high-order interactions
  - Hard to interpret: “The extent to which the effect of  $x_1$  depends upon the value of  $x_2$  depends upon the value of  $x_3$ , dude.”
  - Hard to estimate: need lots of data, must regularize
  - But you might really need them, because conditionality runs deep



The Dude abides high-order interactions



# Higher order interactions

- `data(Wines2012)`
- Judgment of Princeton, 2012
  - New Jersey wines vs fine French wines
- Outcome variable: score
- Predictors:
  - region (NJ/FR)
  - nationality of judge (USA/FR-BE)
  - flight (red/white)



# Higher order interactions

- Predictors: region, nationality of judge, flight
- Consider interactions:
  - Interaction of **region** and **judge** is bias.  
Bias depends upon **flight**.
  - Interaction of **judge** and **flight** is preference.  
Preference depends upon **region**.
  - Interaction of **region** and **flight** is comparative advantage.  
Advantage depends upon **judge**.



# Interaction everywhere

- Interaction, regularization, responsibility
- Next time: Markov Chain Monte Carlo

